



Early Journal Content on JSTOR, Free to Anyone in the World


This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.



XIII. *An Account of some farther Experiments touching the Ascent of Water between two Glass Planes in an Hyperbolick Curve.* By the late Mr. Fr. Hauksbee, F. R. S.

THE Figure of the Hyperbolick Curve, form'd by the Ascending of Water between two Square Glass Planes, as related in *Philosoph. Transact.* Numb. 336. gave me occasion to make some farther Enquiries; and by many Experiments I find, that the same Curve holds in all Directions of the Planes, the Assymtotes being always, one the Surface of the Water, the other a Line drawn along the touching Sides. Thus, when the touching Sides were plung'd under the Surface of the Water, and the Angle c was depress'd and made to remain lower than the Angle a . as in *Fig. 1.* then would be produc'd such an oblique Curve, as may be observ'd in that Figure. In all the several Schemes, ab represents the Surface of the Water on the outside of the Planes; and ac the touching Sides of the same. Now, tho' the Curve dd rises between the Planes in such an obliquity, yet does it conform it self in its Figure to the Assymtotes, viz. ab the Surface of the Water, and ac the touching Sides of the Planes; for supposing the Assymtote ac to be continued, as in the prick'd Line, till it surmounts the Surface of the Water to such a height, or suppose the Planes extended in the same manner, then would the Water remain between them in the appearance of the prick'd Lines, being at all distances from the Axis of the Curve, equal in respect to the Assym-

See Tab. III.

totes; and so of all the rest of the Curves, as in the Figures 2, 3, 4, 5, 6. which are the result of the several Angles, made by the touching Sides of the Planes. Now when the touching Sides were plac'd upwards, parallel to the Surface of the Water (as in *Fig 7.*) and plung'd wholly under the same, then upon lifting them up, in the same Position, till the weight of the Water between the Planes over-balance'd the Power of their Attraction, two Curves, one from each side of the Planes, would open themselves, and meet each other in the middle, as represented in the aforesaid Figure; where they would unite, and make a Figure as join'd by the prick'd Lines, being wider in the middle than towards the sides of the Planes. And it is highly remarkable, that this Curve would always break out between the Planes, at an equal distance between the touching sides and the Surface of the Water.

The same Figure is likewise produc'd between two round Glass Planes, (see *Fig 8.*) the Asymptotes being the same as the former; that is to say, one the Surface of the Water, the other a Tangent drawn from the touching Point, parallel to a Tangent drawn from the open or opposite part of the Planes, being at right Angles with a Line drawn thro' the Axis of the same. These Experiments I find to answer the same in *Vacuo*, as in the open Air; so that that Element has nothing to do in this extraordinary appearance.

The Planes made use of in the foregoing Experiments were about 7 Inches square, open'd on one side to an Angle of 20 Minutes or thereabouts; the round Planes were in Diameter near 3 Inches.